

Recombinant Human PGD2 Synthase/PTGDS Protein (His Tag)

Catalog No. PKSH032006

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Prostaglandin D Synthase;Prostaglandin-H2 D-Isomerase;Beta-Trace Protein;Cerebrin-28;Glutathione-Independent PGD Synthase;Lipocalin-Type Prostaglandin-D Synthase;Prostaglandin-D2 Synthase;PGD2 Synthase;PGDS;PGDS2;PTGDS;PDS
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Gln190
Accession	P41222
Calculated Molecular Weight	20.1 kDa
Observed molecular weight	28 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 80 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 80 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

PTGDS; also known as L-PGDS; belongs to the calycin superfamily; lipocalin family. Lipocalins share limited regions of sequence homology and a common tertiary structure architecture. They transport small hydrophobic molecules such as steroids; bilins; retinoids; and lipids. PTGDS is a glutathione-independent prostaglandin D synthase that catalyzes the conversion of PGH₂ to PGD₂. It is involved in smooth muscle contraction/relaxation and a variety of central nervous system functions. PTGDS may have an anti-apoptotic role in oligodendrocytes. It binds small non-substrate lipophilic molecules; including biliverdin; bilirubin; retinal; retinoic acid and thyroid hormone; and may act as a scavenger for harmful hydrophobic molecules and as a secretory retinoid and thyroid hormone transporter. It is likely to play important roles in both maturation and maintenance of the central nervous system and male reproductive system.

For Research Use Only