A Reliable Research Partner in Life Science and Medicine

Recombinant Human EphA4 Protein

Catalog No. PKSH032008

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Ephrin type-A receptor 4;HEK8;SEK;TYRO1;EPHA4;Tyrosine-protein kinase

receptor SEK;Tyrosine-protein kinase TYRO1;EK8;hEK8;EPH-like kinase 8

Species Human

Expression Host HEK293 Cells
Sequence Val20-Thr547
Accession P54764
Calculated Molecular Weight 59.2 kDa
Observed molecular weight 60-80 kDa

Bioactivity Not validated for activity

Properties

Tag

Purity > 95 % as determined by reducing SDS-PAGE.

None

Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



>95~% as determined by reducing SDS-PAGE.

Background

Ephrin type-A receptor 4(EPHA4) belongs to the protein kinase superfamily and Ephrin receptor subfamily. EPHA4

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

contains 1 Eph LBD domain; 2 fibronectin type-III domains; 1 protein kinase domain and 1 SAM domain. EPH and EPHrelated receptors have been implicated in mediating developmental events; particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com