

## Recombinant Human Activin Receptor 2B/ACVR2B Protein (His Tag)

**Catalog No.** PKSH032040

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Activin Receptor Type-2B;Activin Receptor Type IIB;ACTR-IIB;ACVR2B;Bone Morphogenetic Protein Receptor Type-2;BMP Type-2 Receptor;BMPR-2;Bone Morphogenetic Protein Receptor Type II;BMP Type II Receptor
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Ser19-Thr134
<b>Accession</b>	Q13705
<b>Calculated Molecular Weight</b>	14.4 kDa
<b>Observed molecular weight</b>	25-38 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Activin proteins that belong to the transforming growth factor-beta (TGF- $\beta$ ) superfamily; exert their biological actions by binding to heteromeric receptor complexes of type I and type II serine/threonine kinase receptors. On ligand binding; type I and II receptors form a stable complex; resulting in phosphorylation of type I receptors by type II receptors with constitutive kinase activity; and subsequently initiates the activation of downstream molecules including the endogenous Smads. ActRIIB; also known as ActRIIB; is a type II receptor containing an extracellular domain (ECD); a transmembrane segment; and a cytoplasmic region that includes the kinase domain. ActRIIB is a receptor for activin A; activin B and inhibin A. Multiple ActRIIB isoforms can also be generated; which bind activin isoforms with different affinities.