

# Recombinant Human Astrocytic Phosphoprotein PEA-15/PEA15 Protein

Catalog No. PKSH032094

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Synonyms** Astrocytic Phosphoprotein PEA-15;15 kDa Phosphoprotein Enriched in

Astrocytes; Phosphoprotein Enriched in Diabetes; PED; PEA15

Species Human Expression Host E.coli

Sequence Met 1-Ala130

Accession Q15121
Calculated Molecular Weight 15.3 kDa
Observed molecular weight 12-16 kDa
Tag None

**Bioactivity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs. **Formulation** Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



>95~% as determined by reducing SDS-PAGE.

### **Background**

Astrocyticphosphoprotein PEA-15 (PEA15) is a death effector domain (DED)-containing protein. PEA15 is mainly

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Email: techsupport@elabscience.com

Web: www.elabscience.com





A Reliable Research Partner in Life Science and Medicine

expressed in the central nervous system, principally in astrocytes. Increased PEA15 levels affect tumorigenesis and cancer progression. PEA15 is overexpressed in breast cancers and gliomas as well as in type 2 diabetes. PEA15 blocks Rasmediated inhibition of integrin activation and modulates the ERK MAP kinase cascade. PEA15 also inhibits RPS6KA3 activities by holding it in the cytoplasm. In addition, PEA15 inhibits both TNFRSF6 and TNFRSF1A mediated CASP8 activity and apoptosis. At present, PEA15 expression is also a significant prognostic marker in ovarian cancer.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com