A Reliable Research Partner in Life Science and Medicine

## Recombinant Human BCAS2/DAM1 Protein (His & T7 Tag)

Catalog No. PKSH032111

Note: Centrifuge before opening to ensure complete recovery of vial contents.

## **Description**

Synonyms Pre-mRNA-Splicing Factor SPF27;Breast Carcinoma-Amplified Sequence 2;DNA

Amplified in Mammary Carcinoma 1 Protein; Spliceosome-Associated Protein SPF

27;BCAS2;DAM1

Species Human
Expression Host E.coli

SequenceAla2-Phe225AccessionO75934Calculated Molecular Weight28.6 kDaObserved molecular weight60 kDa

Tag N-T7 & C-His

**Bioactivity** Not validated for activity

#### **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 200mM NaCl,

2mM DTT, pH 8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 90 % as determined by reducing SDS-PAGE.

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

# **Background**

Breast Carcinoma-Amplified Sequence 2 (BCAS2) is a member of the SPF27 family. BCAS2 is a nuclear protein and widely expressed in many rtissues. BCAS2 is identified as being overexpressed in various breast cancer cell lines. BCAS2 is a component of the spliceosome, taking part in the removal of introns from mRNA precursors. BCAS2 interacts with estrogen receptor alpha and beta, thyroid hormone receptor beta, peroxisome proliferator-activated receptor gamma. BCAS2 functions as an ER co-activator and is capable of enhancing ER-mediated transcription.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com