Recombinant Human Caspase-10/CASP10 Protein (His Tag)

Catalog No. PKSH032176

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Caspase-10;CASP-10;Apoptotic Protease Mch-4;FAS-Associated Death Domain Protein Interleukin-1B-Converting Enzyme 2;FLICE2;ICE-Like Apoptotic Protease 4;CASP10;MCH4
Species	Human
Expression Host	E.coli
Sequence	Val220-Ile480
Accession	Q92851-4
Calculated Molecular Weight	30.1 kDa
Observed molecular weight	33 kDa
Tag	C-His
Bioactivity	Not validated for activity
Properties	
Purity	> 70 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.
Formulation	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 8% Sucrose, 1mM DTT, 0.05% Tween80, pH8.5.
Reconstitution	Not Applicable
Data	



> 70 % as determined by reducing SDS-PAGE.

Background

Caspase-10 (CASP10) is a 521 amino acid protein member of the Cysteine-Aspartic Acid Protease (Caspase) family. CASP10 contains two DED (Death Effector) domains and is detectable in most tissues. CASP10 cleavage by Granzyme B and autocatalytic activity generate the two active subunits: Caspase-10 subunit p23/17, Caspase-10 subunit p12. Caspases

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

are a family of cytosolic aspartate-specific cysteine proteases involved in the execution-phase of cell apoptosis, the initiation and execution. Human caspases can be subdivided into three functional groups: cytokine activation (caspase-1, -4, -5, and -13), apoptosis initiation (caspase-2, -8, -9, -and -10), and apoptosis execution (caspase-3, -6, and -7). CASP10 cleaves and activates caspases 3 and 7, but itself is processed by caspase 8. Defects in CASP10 are associated with apoptosis defects seen in type II autoimmune lymphoproliferative syndrome.

For Research Use Only