

## Recombinant Human CCL3 Protein

**Catalog No.** PKSH032197

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	C-C Motif Chemokine 3;G0/G1 Switch Regulatory Protein 19-1;Macrophage Inflammatory Protein 1-Alpha;MIP-1-Alpha;PAT 464.1;SIS-Beta;Small-Inducible Cytokine A3;Tonsillar Lymphocyte LD78 Alpha Protein;CCL3;G0S19-1;MIP1A;SCYA3
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Ala 27-Ala 92
<b>Accession</b>	P10147
<b>Calculated Molecular Weight</b>	8.4 kDa
<b>Observed molecular weight</b>	8-11 kDa
<b>Tag</b>	None
<b>Bioactivity</b>	Measure by its ability to chemoattract human PBMC using a concentration range of 5 -50 ng/mL.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 0.1 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS,pH 8.5. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### For Research Use Only

## Background

Human Chemokine (C-C Motif) Ligand 3 (CCL3) is a small cytokine belonging to the CC chemokine family. CCL3 is primarily expressed in T cells, B cells, and monocytes after antigen or mitogen stimulation. CCL3 exhibits chemoattractive and adhesive effects on lymphocytes. CCL3 exerts multiple effects on hematopoietic precursor cells and inhibits the proliferation of hematopoietic stem cells in vitro as well as in vivo. CCR1 and CCR5 have been identified as functional receptors for CCL3.

## For Research Use Only