Recombinant Human CCL3 Protein

Catalog Number: PKSH032197



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	C-C Motif Chemokine 3;G0/G1 Switch Regulatory Protein 19-1;Macrophage Inflammatory Protein 1-Alpha;MIP-1-Alpha;PAT 464.1;SIS-Beta;Small-Inducible Cytokine A3;Tonsillar Lymphocyte LD78 Alpha Protein;CCL3;G0S19-1;MIP1A;SCYA3	
Species	Human	
Expression Host	E.coli	
Sequence	Ala 27-Ala 92	
Accession	P10147	
Calculated Molecular Weight	8.4 kDa	
Observed molecular weight	8-11 kDa	
Tag	None	
Bioactivity	Measure by its ability to chemoattract human PBMC using a concentration range of 5 -50 ng/mL.	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS,pH 8.5. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	

Data

kDa	MK	R
120		STOCKET
90		
60	A REPORT AND	
40	Cold Street of Cold	Salar and
30	-	
20	-	
14	-	-
	1. 3. 6.	Contraction of the
	111111	TRANSFER
WWW.6	labscien	ce.com

> 95 % as determined by reducing SDS-PAGE.

Background

Human Chemokine (C-C Motif) Ligand 3 (CCL3) is a small cytokine belonging to the CC chemokine family. CCL3 is primarily expressed in T cells, B cells, and monocytes after antigen or mitogen stimulation. CCL3 exhibits chemoattractive and adhesive effects on lymphocytes. CCL3 exerts multiple effects on hematopoietic precursor cells and

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Recombinant Human CCL3 Protein

Catalog Number:PKSH032197



inhibits the proliferation of hematopoietic stem cells in vitro as well as in vivo. CCR1 and CCR5 have been identified as functional receptors for CCL3.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com