A Reliable Research Partner in Life Science and Medicine

Recombinant Human CD79B/B29 Protein (His Tag)

Catalog No. PKSH032224

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms B-Cell Antigen Receptor Complex-Associated Protein Beta Chain;B-Cell-Specific

Glycoprotein B29;Ig-Beta;Immunoglobulin-Associated B29

Protein;CD79b;CD79B;B29;IGB

Species Human

Expression Host HEK293 Cells
Sequence Ala29-Asp159
Accession P40259
Calculated Molecular Weight 16.3 kDa
Observed molecular weight 25-36 kDa

Bioactivity Not validated for activity

Properties

Tag

Purity > 95 % as determined by reducing SDS-PAGE.

C-His

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

CD79B is a single-pass type I membrane protein. CD79B contains one Ig-like V-type domain and one ITAM domain. CD79B is required in cooperation with CD79A for initiation of the signal transduction cascade activated by the B-cell antigen receptor complex (BCR), which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79B enhances phosphorylation of CD79A, possibly by recruiting kinases that phosphorylate CD79A or by recruiting proteins that bind to CD79A and protect it from dephosphorylation.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com