Recombinant Human CFHR5 Protein (His Tag)

Catalog No. PKSH032274

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	Complement factor H-related protein 5;CFHR5;CFHL5;FHR5	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Glu19-Glu569	
Accession	Q9BXR6	
Calculated Molecular Weight	63.5 kDa	
Observed molecular weight	61 kDa	
Tag	C-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB,150mM NaCl,pH7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

kDa	МК	R
170 130		
95		
72		
55		
43	-	
34		
26	Name of	

> 95 % as determined by reducing SDS-PAGE.

Background

Complement factor H-related protein 5(FHR-5 for short), is a secreted protein which contains 9 Sushi (CCP/SCR) domains. It is expressed by the liver and secreted in plasma. The pattern of the deposits is similar to other complement

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

components, suggesting that FHR-5 may play a role in complement activation and regulation. Defects in CFHR5 have been found in patients with atypical hemolytic uremic syndrome and may contribute to the disease. In contrast to typical hemolytic uremic syndrome, atypical forms have a poorer prognosis, with higher death rates and frequent progression to end-stage renal disease.

For Research Use Only