## Recombinant Human Cornulin/CRNN Protein (His Tag)

### Catalog No. PKSH032281

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Cornulin;53 kDa Putative Calcium-Binding Protein;53 kDa Squamous Epithelial- Induced Stress Protein;58 kDa Heat Shock Protein;Squamous Epithelial Heat Shock Protein 53;Tumor-Related Protein;CRNN;C1orf10;DRC1;PDRC1;SEP53
Species	Human
Expression Host	E.coli
Sequence	Met 1-Ser140
Accession	Q9UBG3
Calculated Molecular Weight	17.5 kDa
Observed molecular weight	17 kDa
Tag	N-His
Bioactivity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 95 % as determined by reducing SDS-PAGE.

## Background

#### **For Research Use Only**

# **Elabscience**®

Cornulin is a member of the fused gene family of molecular chaperones. Human Cornulin contains N-terminus EF-hand domains and Ca2+ binding domains, and two glutamine- and threonine-rich 60 amino acid repeats in its C-terminus. Cornulin involves in the mucosal/epithelial immune response and epidermal differentiation. Cornulin is a survival factor that participates in the clonogenicity of squamous esophageal epithelium cell lines, attenuates deoxycholic acid (DCA)-induced apoptotic cell death and release of calcium. When Cornulin is overexpressed in oral squamous carcinoma cell lines, it regulates negatively cell proliferation by the induction of G1 arrest.

For Research Use Only