## Recombinant Human CRTAM /CD355 Protein (His Tag)

Catalog Number: PKSH032286



Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

**Synonyms** Cytotoxic and Regulatory T-Cell Molecule; Class-I MHC-Restricted T-Cell-

Associated Molecule; CD355; CRTAM

**Species** Human

**Expression Host** HEK293 Cells **Sequence** Ser18-Ser286 Accession O95727 Calculated Molecular Weight 31.0 kDa Observed molecular weight 64 kDa C-His Tag

## **Properties**

> 95 % as determined by reducing SDS-PAGE. **Purity** 

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to Storage

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs. **Formulation** 

Lyophilized from a  $0.2~\mu m$  filtered solution of 20mM PB, 150mM NaCl, pH 7.2.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed man

Reconstitution Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Cytotoxic and Regulatory T-Cell Molecule (CRTAM) is a member of Nectin family under the immunoglobulin superfamily that is expressed by activated CD8+ and NK T cells. CRTAM is found in spleen, thymus, small intestine, peripheral blood, and it is highly expressed by Purkinje cells of the cerebellum. CRTAM is a type I transmembrane glycoprotein containing one Ig-like C2-type domain and one Ig-like V-type domain in its extracellular domain, while its cytoplasmic region shows a potential class I PDZ domain. CRTAM is expressed as a homodimer on the cell surface but does not show homotypic binding in trans. The high affinity of CRTAM/IGSF4 adhesion allows CRTAM to disrupt IGSF4 homotypic interactions. IGSF4 and T cell receptor coengagement of CD8+ cells expressiong CRTAM induces

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Email: techsupport@elabscience.com Web: www.elabscience.com

# Recombinant Human CRTAM /CD355 Protein (His Tag)

Catalog Number:PKSH032286



increased IFN $\gamma$  or IL-22 production.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>