

Recombinant Human CXCL7/NAP-2 Protein (His Tag)

Catalog No. PKSH032306

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Platelet Basic Protein;PBP;C-X-C Motif Chemokine 7;Leukocyte-Derived Growth Factor;LDGF;Macrophage-Derived Growth Factor;MDGFSmall-Inducible Cytokine B7;PPBP;CTAP3;CXCL7;SCYB7;TGB1;THBGB1
Species	Human
Expression Host	HEK293 Cells
Sequence	Ser35-Asp128
Accession	P02775
Calculated Molecular Weight	11.3 kDa
Observed molecular weight	12-16 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM HAc-Nac, 150mM NaCl, pH 4.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Human Chemokine (C-X-C motif) Ligand 7 (CXCL7); also known as neutrophil activating peptide 2 (NAP-2); is a member of the CXC chemokines containing an ELR domain (Glu-Leu-Arg tripeptide motif). Similar to other ELR domain containing CXC chemokines; such as IL-8 and the GRO proteins; CXCL7 binds CXCR2; chemoattracts and activates neutrophils. CXCL7; Connective Tissue Activating Protein III (CTAPIII) and β thromboglobulin (β TG); are proteolytically processed carboxylterminal fragments of platelet basic protein (PBP) which is found in the alphagranules of human platelets. Although CTAPIII; β TG; and PBP represent amino-terminal extended variants of NAP2 and possess the same CXC chemokine domains; these proteins do not exhibit CXCL7/NAP2 activity. CXCL7 induces cell migration through the G-protein-linked receptor CXCR-2.

For Research Use Only