Recombinant Human Cyclophilin E/PPIE Protein (His Tag)

Catalog Number: PKSH032317



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Peptidyl-Prolyl Cis-Trans Isomerase E;PPIase E;Cyclophilin

E;Cyclophilin-33;Rotamase E;PPIE;CYP33

SpeciesHumanExpression HostE.coli

Sequence Met 1-Val301
Accession Q9UNP9
Calculated Molecular Weight 35.6 kDa
Observed molecular weight 34 kDa
Tag N-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue

ice/gel packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, pH 8.0.

Reconstitution Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Peptidyl-prolyl cis-trans isomerase E, also known as Cyclophilin E, Cyclophilin-33, Rotamase E, CYP33, PPIE, is an enzyme which belongs to the cyclophilin-type PPIase family of PPIase E subfamily. PPIE found in all the examined tissues including heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. PPIE contains one PPIase cyclophilin-type domain and one RRM (RNA recognition motif) domain. PPIE accelerates the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. PPIE combines RNA-binding and PPIase activities. It may be involved in muscle- and brain-specific processes and pre-mRNA splici

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com