A Reliable Research Partner in Life Science and Medicine

Recombinant Human EGR1 Protein (His Tag)

Catalog No. PKSH032369

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms EGR-1; Early growth response protein 1; Zif 268; zinc finger protein 225; NGFI-

A;nerve growth factor-induced protein A;

SpeciesHumanExpression HostE.coli

Sequence Gln282-Ser433

AccessionP18146Calculated Molecular Weight19.9 kDaObserved molecular weight19 kDaTagN-His

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

EGR-1 belongs to the EGR family of C2H2-type zinc finger proteins. It is a nuclear protein and functions as a

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

transcriptional regulator. EGR-1 recognizes and binds to the DNA sequence 5'-CGCCCCGC-3'(EGR-site). The products of target genes it activates are required for differentiation and mitogenesis. Studies suggest this is a tumor suppressor gene. EGR-1 has a distinct pattern of expression in the brain, and its induction has been shown to be associated with neuronal activity. Several studies suggest it has a role in neuronal plasticity. EGR-1 has also been found to regulate the expression of synaptobrevin II (a protein important for synaptic exocytosis).

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com