

Recombinant Human 4E-BP1/EIF4EBP1 Protein (His Tag)

Catalog No. PKSH032373

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Eukaryotic Translation Initiation Factor 4E-Binding Protein 1;4E-BP1;eIF4E-Binding Protein 1;Phosphorylated Heat- and Acid-Stable Protein Regulated by Insulin 1;PHAS-I;EIF4EBP1;4E-BP1;4EBP1;BP-1
Species	Human
Expression Host	E.coli
Sequence	Met 1-Ile118
Accession	Q13541
Calculated Molecular Weight	14.7 kDa
Observed molecular weight	17 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Eukaryotic Translation Initiation Factor 4E-Binding Protein 1 (4EBP1) is a member of the eIF4E-binding protein family. 4EBP1 regulates eIF4E activity by preventing its assembly into the eIF4F complex. 4EBP1 mediates the regulation of protein translation by hormones, growth factors and other stimuli that signal through the MAP kinase and mTORC1 pathways. Non-phosphorylated 4EBP1 competes with EIF4G1/EIF4G3 to interact with EIF4E. 4EBP1 is phosphorylated in response to various signals including insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation. 4EBP1 has a role in progression of breast neoplasms through cell signaling.