A Reliable Research Partner in Life Science and Medicine

Recombinant Human Ephrin-A4/EFNA4 Protein (His Tag)

Catalog No. PKSH032391

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Ephrin-A4;EPH-Related Receptor Tyrosine Kinase Ligand

4;LERK-4;EFNA4;EPLG4;LERK4

Species Human

Expression Host HEK293 Cells **Sequence** Leu26-Gly171 P52798 Accession Calculated Molecular Weight 17.4 kDa

Observed molecular weight 20 kDa Tag C-His

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to **Storage**

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS5% trehalose, 5% mannitol,

0.01% Tween 80, pH7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Ephrin-A4 is a member of the ephrin ligand family which binds members of the Eph receptor family. All ligands share a conserved extracellular sequence; which most likely corresponds to the receptor binding domain. Ephrin-A4 consists of approximately 125 amino acids and includes four invariant cysteines; It has been shown to bind EphA2; EphA3; EphA4; EphA5; EphA6; EphA7; and EphB1. Ephrin-A4 binds promiscuously Eph receptors residing on adjacent cells; leading to contact-dependent bidirectional signaling into neighboring cells. It may play a role in the interaction between activated Blymphocytes and dendritic cells in tonsils.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com