A Reliable Research Partner in Life Science and Medicine

Recombinant Human FGF-1/FGFa Protein

Catalog No. PKSH032431

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Fibroblast Growth Factor 1;FGF-1;Acidic Fibroblast Growth

Factor;aFGF;Endothelial Cell Growth Factor;ECGFHeparin-Binding Growth Factor

1;HBGF-1;FGF1;FGFA

Species Human
Expression Host E.coli

Sequence Phe16-Asp155

AccessionP05230Calculated Molecular Weight16.8 kDaObserved molecular weight18 kDaTagC-His

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS,pH 8.0.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

FGF acidic; also known as ECGF; FGF-1and HBGF-1; is a non-glycosylated heparin binding growth factor that is

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

expressed in the brain; kidney; retina; smooth muscle cells; bone matrix; osteoblasts; astrocytes and endothelial cells. It is a mitogenic peptide that is produced by multiple cell types and stimulates the proliferation of cells of mesodermal; ectodermal; and endodermal origin. Its association with heparan sulfate is a prerequisite for activation of FGF receptors. Internalized FGF acidic migrates to the nucleus where it is phosphorylated by nuclear PKC delta; exported to the cytosol; dephosphorylated; and degraded. Intracellular FGF acidic inhibits p53 activity and proapoptotic signaling.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com