
Recombinant Human FGF-19/FGF9 Protein (His Tag)

Catalog No. PKSH032436

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|-----------------------------|---|
| Synonyms | Fibroblast growth factor 19;FGF-19;FGF19 |
| Species | Human |
| Expression Host | E.coli |
| Sequence | Phe27-Lys216 |
| Accession | O95750 |
| Calculated Molecular Weight | 22.6 kDa |
| Observed molecular weight | 24 kDa |
| Tag | C-His |
| Bioactivity | Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is < 51 ng/mL. |
| Properties | |
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 0.1 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS,pH 8.0. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |
| Data | |



>95 % as determined by reducing SDS-PAGE.

Background

Fibroblast growth factor 19 (FGF19) is a secreted protein which belongs to the FGFs family. FGF19 is expressed in fetal

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

brain, cartilage, retina, and adult gall bladder. FGFs modulate cellular activity via at least 5 distinct subfamilies of highaffinity FGF receptors (FGFRs): FGFR-1, -2, -3, and -4, all with intrinsic tyrosine kinase activity. FGFRs can be important for regulation of glucose and lipid homeostasis. FGF19 has important roles as a hormone produced in the ileum in response to bile acid absorption. It has been shown to cause resistance to diet-induced obesity and insulin desensitization and to improve insulin, glucose, and lipid profiles in diabetic rodents. FGF19 can be considered as a regulator of energy expenditure.

For Research Use Only