A Reliable Research Partner in Life Science and Medicine

Recombinant Human FGF-2/FGFb Protein

Catalog No. PKSH032438

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Fibroblast growth factor 2;FGF-2;Basic fibroblast growth factor;Bfgf;Heparin-

binding growth factor 2;HBGF-2;FGF2;FGFB

Species Human Expression Host E.coli

Sequence Ala 135-Ser 288

AccessionP09038Calculated Molecular Weight18.1 kDaObserved molecular weight17 kDaTagN-His

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per μ g of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS with 0.01% sarkosyl,pH 8.0.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Fibroblast growth factor 2(FGF2) is a secreted protein and belongs to the heparin-binding growth factors family. FGF2 is produced by epithelial; tumor and other cell types. It involved in developmental processes and regulates differentiation;

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Email: techsupport@elabscience.com

Web: www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

proliferation; and migration; FGF2 is a critical factor for growing embryonic stem cells in culture without inducing differentiation. FGF2 has a high affinity for heparan sulfate and binding is a step in the FGF basic activation of FGFR tyrosine kinase.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com