# Recombinant Human FGF21 Protein (His Tag)

Catalog Number:PKSH032440



Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Synonyms** Fibroblast Growth Factor 21;FGF-21;FGF21

Species Human Expression Host E.coli

SequenceHis29-Ser209AccessionQ9NSA1Calculated Molecular Weight20.4 kDaObserved molecular weight25 kDaTagC-His

**Bioactivity** Measure by its ability to induce proliferation in BaF3 cells transfected with human

FGFRIIIc.The ED<sub>50</sub> for this effect is  $< 0.4 \mu g/mL$ .

#### **Properties**

**Purity** > 98 % as determined by reducing SDS-PAGE.

**Endotoxin** < 0.1 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS,pH 8.0.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 98 % as determined by reducing SDS-PAGE.

#### **Background**

Fibroblast Growth Factor 21 (FGF21) is a growth factor that belongs to the FGF family. FGF family proteins play a central role during prenatal development and postnatal growth and regeneration of mamy tissues, by promoting cellular proliferation and differentiation. FGF21 is a potent activator of glucose uptake on adipocytes, protects animal from dietinduced obesity when overexpression in transgenic mice, and lower blood glucose and triglyceride levels when therapeutically adiministered to diabetic redents. FGF21 is produced by hepatocytes in reponse to free fatty acid stimulation of a PPARa/RXR dimeric complex. This situation occurs clinically during starvation, or following the

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

## **Recombinant Human FGF21 Protein (His Tag)**

Catalog Number:PKSH032440



ingestionof a highly-fat/low-carbohydrate diet.Upon FGF21 secretion, white adipose tissue is induced to release free fatty acids from triglyceride stores. Once free fatty acid reach hepatocytes, they are oxidized and reduced to acetyl-CoA. The acetyl-CoA is recombined into 4-carbon ketone bodies, release, and transported to peripheral tissue for TCA processing and energy generation.

### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com