A Reliable Research Partner in Life Science and Medicine

### Recombinant Human Fibronectin/FN Protein

Catalog No. PKSH032450

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms Fibronectin;FN1;CIG;ED-B;FINC;FN;FNZ;GFND;GFND2;LETS;MSF

Species Human
Expression Host E.coli

Sequence Pro1270-Ser1546&Ala1721-Thr2016

AccessionP02751Calculated Molecular Weight62.7 kDaObserved molecular weight60-80 kDaTagNone

**Bioactivity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 0.01 EU per  $\mu g$  of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 12.5 mM Citric acid, 1.25% Sucrose,

0.1% Tween 80, pH 5.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

# <u>Data</u>



> 95 % as determined by reducing SDS-PAGE.

### **Background**

Fibronectin1(FN1) is a secreted protein and contains 12 fibronectin type-I domains; fibronectin type-II domains and 16

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

 $Email: \underline{tech support@elabscience.com}$ 

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

fibronectin type-III domains. Recombinant human fibronectin fragment; is a protein of ~63 kDa containing a central cellbinding domain; a high affinity heparin-binding domain II; and CS1 site within the alternatively spliced III CS region of human fibronectin. Cells bind to a VLA-4 ligand; a CS-I site; and a VLA-5 ligand; a cell attachment domain; and virus vectors binds to a heparin binding domain II; which co-locates the cell and the virus vector on NovoNectin. This process enhances the density of both cells and vectors; and facilitates the gene transduction in the result.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com