

Recombinant Human GFRA1/GDNFRA Protein (Fc Tag)

Catalog No. PKSH032481

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	GDNF Family Receptor Alpha-1;GDNF Receptor Alpha-1;GDNFR-Alpha-1;GFR-Alpha-1;RET Ligand 1;TGF-Beta-Related Neurotrophic Factor Receptor 1;GFRA1;GDNFRA;RETL1;TRNR1
Species	Human
Expression Host	HEK293 Cells
Sequence	Asp25-Lys429
Accession	P56159-2
Calculated Molecular Weight	72.4 kDa
Observed molecular weight	85 kDa
Tag	C-Fc
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Glial Cell Line-Derived Neurotrophic Factor Family Receptor α -1 (GDNFR α 1) is a glycosylphosphatidylinositol (GPI) linked cell surface protein belonging to GDNF-family receptor α subtype which consists of at least four members. GFR α 1 and GFR α 2 are the cognate co-receptor for the neurotrophic factor neurturin mediating the NRTN-induced autophosphorylation and activation of the RET tyrosine kinase receptor. Soluble GFR α s released enzymatically from the cell surface by phosphatidylinositol phospholipase C, as well as recombinantly produced soluble GFR α 1, can also bind with high affinity to GDNF and trigger the activation of Ret tyrosine kinase. Human GFR α 1 shares 93% amino acid identity with mouse GFR α 1. The expression of the various GFR α s are differentially regulated in the central and peripheral nervous system, suggesting complementary roles for the GFR α s in mediating the activities of the GDNF family of neurotrophic factors.