
# Recombinant Human GITRL/TNFSF18 Protein (His Tag)

#### Catalog No. PKSH032485

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Tumor necrosis factor ligand superfamily member 18;TNFSF18;Activation- inducible TNF-related ligand;Glucocorticoid-induced TNF-related ligand;AITRL; GITRL; TL6
Species	Human
Expression Host	Human Cells
Sequence	Glu74-Ser199
Accession	Q9UNG2
Calculated Molecular Weight	15.3 kDa
Observed molecular weight	16 & 18 kDa
Tag	C-6His
Properties	
Purity	> 90 % as determined by reducing SDS-PAGE.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



## Background

TNFSF18 is a single-pass type II membrane protein. It is expressed at high levels in the small intestine, ovary, testis, kidney and endothelial cells. TNFSF18 cytokine binds to TNFRSF18/AITR/GITR. It regulates T-cell responses, and functions as costimulator and lower the threshold for T-cell activation and T-cell proliferation. It is Important for interactions between activated T-lymphocytes and endothelial cells and Promotes leukocyte adhesion to endothelial cells. TNFSF18 mediates activation of NF-kappa-B. As Triggers increased phosphorylation of STAT1 and up-regulates

### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

# **Elabscience**®

expression of VCAM1 and ICAM1. It also regulates migration of monocytes from the splenic reservoir to sites of inflammation.

**For Research Use Only**