A Reliable Research Partner in Life Science and Medicine

Recombinant Human GRB2 Protein (His Tag)

Catalog No. PKSH032512

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Growth Factor Receptor-Bound Protein 2; Adapter Protein GRB2; Protein

Ash;SH2/SH3 Adapter GRB2;GRB2;ASH

Species Human **Expression Host** E.coli

Sequence Met 1-Val217

P62993 Accession Calculated Molecular Weight 26.3 kDa Observed molecular weight 25-30 kDa Tag C-His

Bioactivity Not validated for activity

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to **Storage**

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM PB, 6% Sucrose, 4%

Mannitol, 50mM NaCl, 0.05% Tween 80, pH8.0.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

As an adaptor protein, Growth Factor Receptor-Bound Protein 2 (GRB2) is involved in siganl transduction and consists of a central SH2 domain flanked by two SH3 domains. GRB2 associates with activated Tyr-phosphorylated EGF receptor/EGFR and PDGF receptors via its SH2 domain, stimulating GTP binding to Ras, which in turn activates MAPK and other signaling pathway. It also associates to other cellular Tyr-phosphorylated proteins such as SIT1, IRS1, IRS4, SHC and LNK. probably via the concerted action of both its SH2 and SH3 domains.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com