
Recombinant Human Haptoglobin/Zonulin Protein (His Tag)

Catalog No. PKSH032517

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Haptoglobin;Zonulin;HP
Species	Human
Expression Host	HEK293 Cells
Sequence	Val19-Gln160(alphachain)&Ile162-Asn406(betachain)
Accession	P00738
Calculated Molecular Weight	15.9&28.3 kDa
Observed molecular weight	16&40-75 kDa
Tag	C-His
Bioactivity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Haptoglobin(HP) is a secreted protein which is a member of the peptidase S1 family of serine proteases. As a result of hemolysis; hemoglobin is found to accumulate in the kidney and is secreted in the urine. Haptoglobin captures; and

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

combines with free plasma hemoglobin to allow hepatic recycling of heme iron and to prevent kidney damage. Haptoglobin also acts as an Antimicrobial; Antioxidant has antibacterial activity and plays a role in modulating many aspects of the acute phase response. Hemoglobin/haptoglobin complexes are rapidely cleared by the macrophage CD163 scavenger receptor expressed on the surface of liver Kupfer cells through an endocytic lysosomal degradation pathway. Uncleaved haptoglogin; also known as zonulin; plays a role in intestinal permeability; allowing intercellular tight junction disassembly; and controlling the equilibrium between tolerance and immunity to non-self antigens.

For Research Use Only