
## Recombinant Human IA2/PTPRN Protein (aa 687-979, His Tag)

#### Catalog No. PKSH032556

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Receptor-type tyrosine-protein phosphatase-like N;R-PTP-N;Islet cell antigen 512;ICA 512;Islet cell autoantigen 3;PTP IA-2;PTPRN;ICA3;ICA512
Species	Human
Expression Host	E.coli
Sequence	Met687-Gln979
Accession	Q16849
Calculated Molecular Weight	35.6 kDa
Observed molecular weight	33 kDa
Tag	N-His
Bioactivity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.
Storage	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.
Formulation	Supplied as a 0.2 $\mu$ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
Reconstitution	Not Applicable
Data	



> 95 % as determined by reducing SDS-PAGE.

### Background

Receptor-type tyrosine-protein phosphatase-like N (PTPRN) belongs to the protein-tyrosine phosphatase family and receptor class 8 subfamily. PTPRN contains 1 tyrosine-protein phosphatase domain; is expressed in neuroendocrine cells only. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth; differentiation; mitotic cycle; and oncogenic transformation. It implicated in neuroendocrine secretory processes. It may be involved in processes specific for neurosecretory granules; such as their biogenesis; trafficking or regulated exocytosis

#### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

# **Elabscience**®

or may have a general role in neuroendocrine functions. It seems to lack intrinsic enzyme activity; may play a role in the regulation of secretory granules via its interaction with SNTB2. This PTP was found to be an autoantigen that is reactive with insulin-dependent diabetes mellitus (IDDM) patient sera; and thus may be a potential target of autoimmunity in diabetes mellitus.

**For Research Use Only**