

Recombinant Human IL2Ra/CD25 Protein (Fc Tag)

Catalog No. PKSH032570

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-2 receptor subunit alpha;CD25;p55;TAC antigen;IL2-RA;IL-2R subunit alpha;IL-2-RA;IL-2 receptor subunit alpha;IL-2Rα;IDDM10;IL2R;p55;TCGFR
Species	Human
Expression Host	HEK293 Cells
Sequence	Glu22-Cys213
Accession	P01589
Calculated Molecular Weight	48.9 kDa
Observed molecular weight	60-80 kDa
Tag	C-Fc
Bioactivity	Loaded Human IL-2-His on HIS1K Biosensor, can bind Human IL-2RA-Fc with an affinity constant of 3.06 nM as determined in BLI assay.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Interleukin-2 receptor subunit alpha (IL2RA) is a single-pass type I membrane protein; contains 2 Sushi (CCP/SCR) domains. The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains; together with the common gamma chain (IL2RG); constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor; while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein; soluble IL2RA has been isolated and determined to result from extracellular proteolysis.