

Recombinant Human IL-20RB Protein (Fc Tag)

Catalog No. PKSH032574

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-20 receptor subunit beta;IL-20 receptor subunit beta;IL-20R-beta;IL-20RB;IL-20R2;DIRS1;hCG_2022374;FNDC6;MGC34923;fibronectin type III domain containing 6;interleukin-20 receptor II
Species	Human
Expression Host	HEK293 Cells
Sequence	Asp30-Ala230
Accession	Q6UXL0
Calculated Molecular Weight	49.6 kDa
Observed molecular weight	60-85 kDa
Tag	C-Fc
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Reconstitution	Not Applicable

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Interleukin-20 receptor subunit beta (IL20RB) is a single-pass type I membrane protein and belongs to the type II cytokine receptor family. It contains 2 fibronectin type-III domains. There are two kinds of type II cytokine receptors : cytokine receptors that bind type I and type II interferons; cytokine receptors that bind members of the interleukin-10 family (interleukin-10, interleukin-20 and interleukin-22). Type II cytokine receptors are similar to type I cytokine

For Research Use Only

receptors except they do not possess the signature sequence WSXWS that is characteristic of type I receptors. They are expressed on the surface of certain cells, which bind and respond to a select group of cytokines. These receptors are related predominantly by sequence similarities in their extracellular portions that are composed of tandem Ig-like domains. The intracellular domain of type II cytokine receptors is typically associated with a tyrosine kinase belonging to the Janus kinase (JAK) family