

Recombinant Human Inhibin β C Chain/INHBC Protein (aa 19-352, His Tag)

Catalog No. PKSH032586

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Inhibin Beta C Chain;Activin Beta-C Chain;INHBC
Species	Human
Expression Host	HEK293 Cells
Sequence	Thr19-Ser352
Accession	P55103
Calculated Molecular Weight	37.5 kDa
Observed molecular weight	42-58&32-40&14-17 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Inhibin beta C chain; also known as activin beta-C chain and INHBC; belongs to the TGF-beta family. INHBC forms a homodimeric or heterodimeric through association with alpha and beta subunits; linked by one or more disulfide bonds.

For Research Use Only

Inhibins are heterodimers of one alpha and one beta subunit. Activins are homo- or heterodimers of beta subunits only. Inhibins/activins regulates many physiological processes; such as hypothalamic and pituitary hormone secretion; gonadal hormone secretion; germ cell development and maturation; erythroid differentiation; insulin secretion; nerve cell survival; embryonic axial development or bone growth and so on.