

# Recombinant Human IDE/Insulysin Protein (His Tag)

Catalog Number:PKSH032593



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                                    |   |
|------------------------------------|---|
| <b>Synonyms</b>                    | Insulin-Degrading Enzyme;Abeta-Degrading Protease;Insulin Protease;Insulinase;Insulysin;IDE |
| <b>Species</b>                     | Human   |
| <b>Expression Host</b>             | HEK293 Cells  |
| <b>Sequence</b>                    | Met42-Leu1019   |
| <b>Accession</b>                   | P14735  |
| <b>Calculated Molecular Weight</b> | 114.3 kDa   |
| <b>Observed molecular weight</b>   | 120 kDa   |
| <b>Tag</b>                         | C-His   |

## Properties

|                       |  |
|-----------------------|--|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.   |
| <b>Endotoxin</b>      | < 1.0 EU per µg of the protein as determined by the LAL method.  |
| <b>Storage</b>        | Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.   |
| <b>Shipping</b>       | This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C. |
| <b>Formulation</b>    | Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 0.05% Brij35, 10% Glycerol, pH 7.5.   |
| <b>Reconstitution</b> | Not Applicable   |

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Insulin-Degrading Enzyme (IDE) is a secreted enzyme that belongs to the peptidase M16 family. IDE is a large zinc-binding protease and cleaves multiple short polypeptides that vary considerably in sequence. IDE plays a role in the cellular breakdown of insulin, IAPP, glucagon, bradykinin, kallidin, and other peptides, and thereby plays a role in intercellular peptide signaling. IDE degrades amyloid formed by APP and IAPP. IDE may participate in the degradation and clearance of naturally secreted amyloid  $\beta$ -protein by neurons and microglia. IDE, which migrates at 110 kDa during gel electrophoresis under denaturing conditions, has since been shown to have additional substrates, including the signaling peptides glucagon, TGF  $\alpha$  and  $\beta$ -endorphin.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017