Recombinant Human IFNAR2 Protein (His Tag)

Catalog Number: PKSH032605



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Interferon Alpha/Beta Receptor 2;IFN-R-2;IFN-Alpha Binding Protein;IFN-

Alpha/Beta Receptor 2;Interferon Alpha Binding Protein;Type I Interferon Receptor

2;IFNAR2;IFNABR;IFNARB

Species Human

Expression Host

Sequence

Ile27-Lys243

Accession

P48551

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

1le27-Lys243

25.8 kDa

38-55 kDa

C-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interferon α/β Receptor 2 (IFN- α/β R2) is a single-pass type I membrane protein which belongs to the type II cytokine receptor family. It complexes with IFN- α/β R1 to form the signaling receptor complex for the family of α and β IFN subtypes. By alternative splicing; IFN- α/β R2 can exist as a secreted soluble protein or as a type I membrane protein. IFN- α/β R2 is the principal ligand binding subunit of the receptor. Ligand binding is stabilized by the subsequent association with IFN- α/β R1; resulting in the formation of a signaling ternary receptor complex. IFNAR2 was detected in most lymphocytes; monocytes; and granulocytes; although IFNAR2 expression was higher in the monocytes and granulocytes

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Recombinant Human IFNAR2 Protein (His Tag)

Catalog Number:PKSH032605



than in the lymphocytes. Among the lymphocyte subsets; IFNAR2 showed high expression in natural killer (NK) cells and low expression in T lymphocytes. Isoform 1 and isoform 3 of IFNAR2 are directly involved in signal transduction due to their interaction with the TYR kinase; JAK1. Isoform 1 also interacts with the transcriptional factors; STAT1 and STAT2. Both forms are potent inhibitors of type I IFN activity.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com