

Recombinant Human IFNAR2 Protein (Fc Tag)

Catalog No. PKSH032606

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interferon Alpha/Beta Receptor 2;IFN-R-2;IFN-Alpha Binding Protein;IFN-Alpha/Beta Receptor 2;Interferon Alpha Binding Protein;Type I Interferon Receptor 2;IFNAR2;IFNABR;IFNARB
Species	Human
Expression Host	HEK293 Cells
Sequence	Ile27-Lys243
Accession	P48551
Calculated Molecular Weight	51.8 kDa
Observed molecular weight	67-75 kDa
Tag	C-Fc
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Interferon α/β Receptor 2 (IFN- α/β R2) is a single-pass type I membrane protein which belongs to the type II cytokine receptor family. It complexes with IFN- α/β R1 to form the signaling receptor complex for the family of α and β IFN subtypes. By alternative splicing; IFN- α/β R2 can exist as a secreted soluble protein or as a type I membrane protein. IFN- α/β R2 is the principal ligand binding subunit of the receptor. Ligand binding is stabilized by the subsequent association with IFN- α/β R1; resulting in the formation of a signaling ternary receptor complex. IFNAR2 was detected in most lymphocytes; monocytes; and granulocytes; although IFNAR2 expression was higher in the monocytes and granulocytes than in the lymphocytes. Among the lymphocyte subsets; IFNAR2 showed high expression in natural killer (NK) cells and low expression in T lymphocytes. Isoform 1 and isoform 3 of IFNAR2 are directly involved in signal transduction due to their interaction with the TYR kinase; JAK1. Isoform 1 also interacts with the transcriptional factors; STAT1 and STAT2. Both forms are potent inhibitors of type I IFN activity.

For Research Use Only