

Recombinant Human Interferon γ /IFNG Protein

Catalog No. PKSH032611

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interferon Gamma, IFN-Gamma, Immune Interferon, IFNG
Species	Human
Expression Host	HEK293 Cells
Sequence	Gln24-Gln166
Accession	P01579
Calculated Molecular Weight	16.8 kDa
Observed molecular weight	16&20-25 kDa
Tag	None
Bioactivity	Testing in progress

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed man
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

IFN γ is the major interferon produced by mitogenically or antigenically stimulated lymphocytes. It is structurally different from type I interferon and its major activity is immunoregulation. It has been implicated in the expression of

For Research Use Only

class II histocompatibility antigens in cells that do not normally produce them; leading to autoimmune disease. Interferon gamma is produced mainly by T-cells and natural killer cells activated by antigens; mitogens; or alloantigens. It is produced by lymphocytes expressing the surface antigens CD4 and CD8. IFN γ synthesis is induced by IL-2; FGF-basic; and EGF.