Recombinant Human Interleukin-17F/IL-17F Protein

Catalog Number: PKSH032623



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | | |
|-----------------------------|--|--|--|
| Synonyms | Interleukin-17F;IL-17F;Cytokine ML-1;Interleukin-24;IL-24;IL17F;IL24 | | |
| Species | Human | | |
| Expression Host | HEK293 Cells | | |
| Sequence | Arg31-Gln163 | | |
| Accession | AAH70124.1 | | |
| Calculated Molecular Weight | 14.9 kDa | | |
| Observed molecular weight | 18 kDa | | |
| Tag | None | | |
| Properties | | | |
| Purity | > 95 % as determined by reducing SDS-PAGE. | | |
| Endotoxin | < 1.0 EU per μ g of the protein as determined by the LAL method. | | |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. | | |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. | | |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed man | | |
| Reconstitution | Please refer to the printed manual for detailed information. | | |
| Data | | | |

| kDa | MK | R | |
|-----------------------|----|---|--|
| 120 90 60 40 | | | |
| 30 | | | |
| 20 | | - | |
| 14 | - | | |
| | | | |

> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-17F (IL-17F) exists in a disulfide-linked heterodimer that belongs to the IL-17 family. IL-17F is expressed in activated; but not resting; CD4+ T-cells and activated monocytes. IL-17F has been shown to stimulate the production of several other cytokines; including IL-6; IL-8; and granulocyte colony-stimulating factor. IL-17F can regulate cartilage matrix turnover and stimulates PBMC and T-cell proliferation. IL-17F is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2; TGFB1/TGFB; and monocyte chemoattractant protein-1. Defects in IL-17F are the cause of familial candidiasis type 6 (CANDF6).

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com