A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL18BP Protein (Fc Tag)

Catalog No. PKSH032626

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Interleukin-18-binding protein; Tadekinig-alfa and IL18BP;

Species Human

Expression Host HEK293 Cells
Sequence Thr31-Gly194
Accession O95998
Calculated Molecular Weight 44.8 kDa
Observed molecular weight 60-75 kDa
Tag C-Fc

Bioactivity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-18-binding protein, also known as Tadekinig-alfa and IL18BP, contains 1 Ig-like C2-type domain. IL18BP is constitutively expressed and secreted in mononuclear cells. IL18BP functions as an IL18 inhibitor. IL18BP binds to IL18,

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Email: techsupport@elabscience.com

Web: www.elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

prevents the binding of IL18 to its receptor, and thus inhibits IL18-induced IFN-gamma production. It has been shown that IL18BP may be a promising molecular approach to inhibit neointimal hyperplasia and arteriosclerosis progression following coronary and peripheral angioplasty.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com