

Recombinant Human Interleukin-2/IL-2 Protein (His Tag)

Catalog No. PKSH032629

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Interleukin-2;IL-2;T-Cell Growth Factor;TCGF;Aldesleukin;IL4
Species	Human
Expression Host	HEK293 Cells
Sequence	Ala21-Thr153
Accession	P60568
Calculated Molecular Weight	16.4 kDa
Observed molecular weight	14-18 kDa
Tag	C-His
Bioactivity	Measured in a cell proliferation assay using CTLL-2 mouse cytotoxic T cells The specific activity of Recombinant Human IL-2 is $\geq 1 \times 10^7$ IU/mg.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.01 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution of 10mM Sodium Citrate, 4% Mannitol, pH4.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Interleukin-2(IL-2) is an interleukin; a type of cytokine signaling molecule in the immune system; belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells; NK cells; lymphokine-activated killer cells; monocytes; macrophages and oligodendrocytes.