

Recombinant Human ITIH3 Protein (His Tag)

Catalog No. PKSH032658

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Inter-alpha-trypsin inhibitor heavy chain H3;ITI heavy chain H3;ITI-HC3;Inter-alpha-inhibitor heavy chain 3;Serum-derived hyaluronan-associated protein;SHAP
Species	Human
Expression Host	HEK293 Cells
Sequence	Leu35-Asp651
Accession	Q06033
Calculated Molecular Weight	70.4 kDa
Observed molecular weight	80-100 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

ITIH3, which is short for Inter-alpha-trypsin inhibitor heavy chain H3, is a 890 aa. protein. It is secreted expression, and

For Research Use Only

belongs to the ITIH family. I-alpha-I plasma protease inhibitors are assembled from one or two heavy chains (H1, H2 or H3) and one light chain, bikunin. Inter-alpha-inhibitor (I-alpha-I) is composed of H1, H2 and bikunin, inter-alpha-like inhibitor (I-alpha-LI) of H2 and bikunin, and pre-alpha-inhibitor (P-alpha-I) of H3 and bikunin. ITTH3 may act as a carrier of hyaluronan in serum or as a binding protein between hyaluronan and other matrix protein, including those on cell surfaces in tissues to regulate the localization, synthesis and degradation of hyaluronan which are essential to cells undergoing biological processes.