

Recombinant Human IPPI2/IDI2 Protein (His Tag)

Catalog No. PKSH032661

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Isopentenyl-Diphosphate Delta-Isomerase 2;Isopentenyl Pyrophosphate Isomerase 2;IPP Isomerase 2;IPPI2;IDI2
Species	Human
Expression Host	E.coli
Sequence	Met 1-Val227
Accession	Q9BXS1
Calculated Molecular Weight	28.9 kDa
Observed molecular weight	28-31 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 1mM DTT, 0.1mM PMSF, pH 8.0.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Isopentenyl Pyrophosphate Isomerase 2 (IDI2) belongs to the IPP isomerase type 1 family. Both isozymes, IDI1 and IDI2 are localized to the peroxisome by a PTS1-dependent pathway. IDI2 is expressed in skeletal muscle, which contains one nudix hydrolase domain. IDI2 binds one magnesium per subunit. IDI2 catalyzes the 1,3-allylic rearrangement of the homoallylic substrate isopentenyl (IPP) to its highly electrophilic allylic isomer, dimethylallyl diphosphate (DMAPP). It is

For Research Use Only

reported that IDI2 is regulated independently from IDI1, by a mechanism that may involve PPAR- α .