Recombinant Human Lactotransferrin/LTF Protein (His

Tag)

Catalog Number: PKSH032679



Note: Centrifuge before opening to ensure complete recovery of vial contents.

T .	.
LIACOPT	ntion
Descri	77717

Synonyms Lactotransferrin; Lactoferrin; Talalactoferrin; Kaliocin-1; Lactoferroxin-

A;Lactoferroxin-B;Lactoferroxin-C;LTF;LF;GIG12;HEL110;HLF2

Species Human

Expression Host

Sequence

Gly20-Lys711

Accession

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Gly20-Lys711

AAH15822.1

77.3 kDa

90 kDa

C-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs. **Formulation** Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Normalla 5 % - 0 % Arabalana manaital and 0.01% Towns 0.0 and did a


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed man

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Lactotransferrin is a member of the transferrin family that transfer iron to the cells and control the level of free iron in the blood and external secretions. Lactotransferrin is a secreted protein and contains two transferrin-like domains. Lactotransferrin can be cleaved into the following four chains: Kaliocin-1; Lactoferroxin-A; Lactoferroxin-B; and Lactoferroxin-C. Lactoferroxin A; Lactoferroxin B; and Lactoferroxin C have opioid antagonist activity. Lactoferroxin A shows preference for mu-receptors; while Lactoferroxin B and Lactoferroxin C have somewhat higher degrees of preference for kappa-receptors than for mu-receptors. LTF has antimicrobial activity (bacteriocide; fungicide) and is part of the innate defense; mainly at mucoses.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com