

Recombinant Human LIF Protein (E.coli)

Catalog No. PKSH032694

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Leukemia Inhibitory Factor;LIF;Differentiation-Stimulating Factor;D Factor;Melanoma-Derived LPL Inhibitor;MLPLI;Emfilermin;LIF;HILDA;CDF;DIA
Species	Human
Expression Host	E.coli
Sequence	Ser23-Phe202
Accession	P15018
Calculated Molecular Weight	20.5 kDa
Observed molecular weight	18 kDa
Tag	N-His
Bioactivity	Measure by its ability to induce TF-1 cells proliferation. The ED ₅₀ for this effect is < 0.2 ng/mL.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 0.1 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

Leukemia Inhibitory Factor (LIF) is a lymphoid factor that promotes long-term maintenance of embryonic stem cells by suppressing spontaneous differentiation. LIF has a number of other activities including cholinergic neuron differentiation; control of stem cell pluripotency; bone and fat metabolism; mitogenesis of certain factor dependent cell lines and promotion of megakaryocyte production in vivo. Human and murine mature LIF exhibit a 78% sequence identity at the amino acid level. Human LIF is equally active on human and mouse cells. Murine LIF is approximately 1000 fold less active on human cells than human LIF.

For Research Use Only