

Recombinant Human CD300a/LMIR1 Protein (His Tag)

Catalog No. PKSH032699

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CMRF35-Like Molecule 8;CLM-8;CD300 Antigen-Like Family Member A;CMRF-35-H9;CMRF35-H9;CMRF35-H;IRC1/IRC2;Immunoglobulin Superfamily Member 12;IgSF12;Inhibitory Receptor Protein 60;IRp60;NK Inhibitory Receptor;CD300a;CD300A;CMRF35H;IGSF12
Species	Human
Expression Host	HEK293 Cells
Sequence	Leu18-Gln178
Accession	Q9UGN4
Calculated Molecular Weight	18.5 kDa
Observed molecular weight	23-37 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.2. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

For Research Use Only

Background

CD300A is a single-pass type I membrane protein that belongs to the CD300 family. CD300A consists of a 163 amino acid (aa) extracellular domain (ECD) with one Ig-like V- type domain; a 21 amino acid transmembrane segment; and a 98 amino acid cytoplasmic domain with tyrosine residues. CD300A is expressed not only by natural killer (NK) cells but also by T-cell subsets; B-cells; dendritic cells; mast cells; granulocytes and monocytes. CD300A is an inhibitory receptor which may contribute to the down-regulation of cytolytic activity in natural killer (NK) cells; and to the down-regulation of mast cell degranulation.