

## Recombinant Human LTBR/TNFRSF3 Protein (His Tag)

Catalog No. PKSH032719

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Tumor Necrosis Factor Receptor Superfamily Member 3;Lymphotoxin-Beta Receptor;Tumor Necrosis Factor C Receptor;Tumor Necrosis Factor Receptor 2-Related Protein;Tumor Necrosis Factor Receptor Type III;TNF-RIII;TNFR-III;LTBR;D12S370;TNFCR;TNFR3;TNFRSF3
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Gln31-Met227
<b>Accession</b>	P36941
<b>Calculated Molecular Weight</b>	22.8 kDa
<b>Observed molecular weight</b>	29-40 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### For Research Use Only

## Background

Tumor necrosis factor receptor superfamily member 3; also known as Lymphotoxin-beta receptor;Tumor necrosis factor C receptor;Tumor necrosis factor receptor 2-related protein;Tumor necrosis factor receptor type III;LTBR;TNFCR; TNFR3 and TNFRSF3; is a member of the tumor necrosis factor (TNF) family of receptors. LTBR is a single-pass type I membrane protein and contains four TNFR-Cys repeats. It is expressed on the surface of most cell types; but not on T and B lymphocytes. LTBR and its ligand play a role in the development and organization of lymphoid tissue and transformed cells. Activation of LTBR can trigger apoptosis. In addition; LTBR can lead to the release of the cytokine interleukin 8.

## For Research Use Only