

Recombinant Human MBIP Protein (His Tag)

Catalog No. PKSH032739

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	MAP3K12-Binding Inhibitory Protein 1;MAPK Upstream Kinase-Binding Inhibitory Protein;MUK-Binding Inhibitory Protein;MBIP
Species	Human
Expression Host	E.coli
Sequence	Met 1-Pro344
Accession	Q9NS73
Calculated Molecular Weight	42.5 kDa
Observed molecular weight	50 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, pH 8.0. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

MAP3K12-binding inhibitory protein 1 (MBIP) is a 39kD protein high expression in the heart and lung. It is a component

For Research Use Only

of the ADA2A-containing complex (ATAC) complex, a complex with histone acetyltransferase activity on histones H3 and H4, and composed of CSRP2BP, KAT2A, TADA2L, TADA3L, ZZ3, MBIP, WDR5, YEATS2, CCDC101 and DR1. In the complex, it probably interacts directly with KAT2A, CSRP2BP and WDR5. It's function to inhibit the MAP3K12 activity to induce the activation of the JNK/SAPK pathway.