

Recombinant Human NGAL/Lipocalin-2 Protein (His Tag, Human Cells)

Catalog No. PKSH032806

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|---|
| Synonyms | Neutrophil gelatinase-associated lipocalin;NGAL;25 kDa alpha-2-microglobulin-related subunit of MMP-9;Lipocalin-2;Oncogene 24p3;Siderocalin LCN2;p25;HNL;NGAL |
| Species | Human |
| Expression Host | HEK293 Cells |
| Sequence | Gln21-Gly198 |
| Accession | P80188 |
| Calculated Molecular Weight | 21.6 kDa |
| Observed molecular weight | 23 kDa |
| Tag | C-His |
| Bioactivity | Not validated for activity |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles. |
| Shipping | This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C. |
| Formulation | Supplied as a 0.2 µm filtered solution of PBS, 50% Glycerol, pH 7.4. |
| Reconstitution | Not Applicable |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

LCN2 is iron-trafficking protein involved in multiple processes such as apoptosis; innate immunity and renal development. LCN2 binds iron through association with 2;5-dihydroxybenzoic acid (2;5-DHBA); a siderophore that shares structural similarities with bacterial enterobactin; and delivers or removes iron from the cell; depending on the context. LCN2 is involved in apoptosis due to interleukin-3 (IL3) deprivation: iron-loaded form increases intracellular

For Research Use Only

iron concentration without promoting apoptosis; while iron-free form decreases intracellular iron levels; inducing expression of the proapoptotic protein BCL2L1/BIM; resulting in apoptosis. LCN2 is involved in innate immunity; possibly by sequestering iron; leading to limit bacterial growth.