

Recombinant Human NGAL/Lipocalin-2 Protein (His Tag, E.coli)

Catalog No. PKSH032807

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Neutrophil gelatinase-associated lipocalin;NGAL;25 kDa alpha-2-microglobulin-related subunit of MMP-9;Lipocalin-2;Oncogene 24p3;Siderocalin LCN2;p25;HNL;NGAL
Species	Human
Expression Host	E.coli
Sequence	Gln21-Gly198
Accession	P80188
Calculated Molecular Weight	21.8 kDa
Observed molecular weight	20-23 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
Formulation	Supplied as a 0.2 µm filtered solution of PBS, 50% Glycerol, pH 7.4.
Reconstitution	Not Applicable

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Neutrophil gelatinase-associated lipocalin(LCN2) is a secreted protein and belongs to the calycin superfamily. This protein is released from injured tubular cells after various damaging stimuli; is already known by nephrologists as one of the most promising biomarkers of incoming Acute Kidney Injury (AKI). Recent evidence also suggests its role as a biomarker in a variety of other renal and non-renal conditions. Moreover; recent studies seem to suggest a potential

For Research Use Only

involvement of this factor also in the genesis and progression of chronic kidney diseases. NGAL is the first known mammalian protein which specifically binds organic molecules called siderophores; which are high-affinity iron chelators. NGAL; first known as an antibacterial factor of natural immunity; and an acute phase protein; is currently one of the most interesting and enigmatic proteins involved in the process of tumor development. acting as an intracellular iron carrier and protecting MMP9 from proteolytic degradation; NGAL has a clear pro-tumoral effect; as has already been observed in different tumors (e.g. breast; stomach; oesophagus; brain) in humans. In thyroid carcinomas; NGAL is strongly induced by NF- κ B; an important factor involved both in tumor growth and in the link between chronic inflammation and neoplastic development. Thus; Lipocalin-2 (LCN2/NGAL) has been implicated in a variety of processes including cell differentiation; proliferation; survival and morphogenesis.