Recombinant Human NKG2DL/ULBP-1 Protein (His Tag)

Catalog Number: PKSH032814



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms NKG2D ligand 1;N2DL-1;NKG2DL1;ALCAN-beta;Retinoic acid early transcript

1I;UL16-binding protein 1;ULBP1

Species Human

Expression Host

Sequence

Gly26-Pro215

Accession

Q9BZM6

Calculated Molecular Weight

Observed molecular weight

Tag

HEK293 Cells

Gly26-Pro215

29BZM6

23.3 kDa

25-30 kDa

C-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

ULBP1; also known as RAET1I and NKG2DL1; is a member of the ULBP/RAET1 gene family. ULBP1 plays an important role in immune responses; especially in cancer and infectious diseases; and is well-known to bind to NKG2D together with at least ULBP 2 and 3. These proteins are distantly related to major histocompatibility class I (MHC I) molecules; possessing the alpha 1 and alpha 2 Ig-like domains; but lacking the alpha 3 domain. Unlike MHC Class I; they have no capacity to bind peptide or interact with beta2-microglobulin. It can activate multiple signaling pathways in primary NK cells; gamma delta T cells; and CD8+ alpha beta T cells; resulting in the production of cytokines and chemokines.ULBP1 is expressed in wide range of tissues including heart; brain; lung; liver; bone marrow and some tumor

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human NKG2DL/ULBP-1 Protein (His Tag)

Catalog Number:PKSH032814



cells; T-cells; B-cells; As an unconventional member of the MHC class I family; ULBP1 is able to interact with soluble CMV glycoprotein UL16 in CMV infected cells. The interaction with UL16 blocked the interaction with the NKG2D receptor; and thus might escape the immune surveillance. Furthermore; UL16 also causes ULBP1 to be retained in the ER and cis-Golgi apparatus so that it does not reach the cell surface. The ULBP1 regulation may have implications for development of new therapeutic strategies against cancer cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com