

Recombinant Human NKG2DL/ULBP-1 Protein (Fc Tag)



Catalog Number:PKSH032815

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|--|
| Synonyms | NKG2D ligand 1;NKG2DL1;ALCAN-beta;Retinoic acid early transcript 1I;UL16-binding protein 1;ULBP1 |
| Species | Human |
| Expression Host | HEK293 Cells |
| Sequence | Gly26-Pro215 |
| Accession | Q9BZM6 |
| Calculated Molecular Weight | 49.4 kDa |
| Observed molecular weight | 58-70 kDa |
| Tag | C-Fc |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

NKG2D ligand 1; also called ULBP1; is a member of UL16-binding protein (ULBP) family which has also been termed the retinoic acid early transcript 1 (RAET1) family. Unlike the classical MHC class I molecules and the MIC molecules possess $\alpha 1$; $\alpha 2$ and $\alpha 3$ domains; ULBP/RAET1 family members lack $\alpha 3$ domain. ULBP1 is recognized by the activating receptor NKG2D on the surface of cytotoxic natural killer (NK) and T cells; and then promotes the lysis of cells expressing ULBP1 which is important for the immune surveillance. ULBP1 and several other family members; ULBP2 and ULBP5; own the ability to bind the human cytomegalovirus (CMV) UL16 glycoprotein. The human CMV glycoprotein UL16 binds to intracellular ULBP1 and so inhibits its expression at the cell surface; which reduces the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human NKG2DL/ULBP-1 Protein (Fc Tag)

Catalog Number:PKSH032815



susceptibility of the virus-infected cell to cytotoxic destruction by NK cells. The expression of ULBP1 has been found on some tumor cells and is implicated in tumor surveillance.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017