Recombinant Human NgR3/RTN4RL1 Protein (His Tag)

Catalog No. PKSH032820

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Reticulon-4 Receptor-Like 1;Nogo Receptor-Like 2;Nogo-66 Receptor Homolog 2;Nogo-66 Receptor-Related Protein 3;NgR3;RTN4RL1;NGRH2;NGRL2
Species	Human
Expression Host	HEK293 Cells
Sequence	Cys25-Ala419
Accession	Q86UN2
Calculated Molecular Weight	45.5 kDa
Observed molecular weight	65 kDa
Tag	C-His
Bioactivity	Not validated for activity
Properties	
Purity	>95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	

kDa MK R 120 90 60 40 30 20 14

> 95 % as determined by reducing SDS-PAGE.

Background

Nogo-66 Receptor-Related Protein 3 (NgR3) has primary structures with NgR2 (NgRH1, NgRL3) and biochemical

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

properties that are homologous to Nogo-66 receptor (NgR), and constitute a novel neuronal receptor protein family. NgR is GPI-anchored and contains eight leucine-rich repeats (LRR), it is the neuronal receptor for the myelin- associated proteins Nogo-A, OMgp (oligodendrocyte myelin glycoprotein), and MAG (myelin-associated glycoprotein) and mediates the inhibition of CNS axonal regeneration both in vitro and in vivo. NgR2 and NgR3 have similar structure and distinct but overlapping expression versus NgR. NgR2 can be metalloproteinase-cleaved to release a soluble ectodomain. NgR2 has also been shown to bind MAG, but ligands for NgR3 have not yet been determined. Mature huaman NgR3 shares 88%, 88%, 48% and 44% amino acid identity with mature mouse NgR3, rat NgR3, human NgRH1 and NgR, repectively.

For Research Use Only