

Recombinant Human OBFC1/STN1 Protein (His Tag)

Catalog No. PKSH032832

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CST Complex Subunit STN1;Oligonucleotide/Oligosaccharide-Binding Fold-Containing Protein 1;Suppressor of Cdc Thirteen Homolog;OBFC1;STN1
Species	Human
Expression Host	E.coli
Sequence	Met 1-Phe368
Accession	AAH17400.1
Calculated Molecular Weight	44.3 kDa
Observed molecular weight	44 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 80 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 1mM DTT, 5% Trehalose, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 80 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

CST Complex Subunit STN1 (OBFC1) is a 368 amino acid protein that contains one OB DNA-binding domain. It is a member of the STN1 family. OBFC1 is component of the CST complex, a complex that binds to single-stranded DNA and is required to protect telomeres from DNA degradation. The CST complex binds single-stranded DNA with high affinity in a sequence-independent manner, while isolated subunits bind DNA with low affinity by themselves. In addition to telomere protection, the CST complex has probably a more general role in DNA metabolism at non-telomeric sites.